



GLOBAL JOURNAL OF MEDICAL RESEARCH: L  
NUTRITION & FOOD SCIENCE  
Volume 22 Issue 3 Version 1.0 Year 2022  
Type: Double Blind Peer Reviewed International Research Journal  
Publisher: Global Journals Inc. (USA)  
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

## Questionnaire Survey Results on Lifestyle Habits (Drinking, Smoking, Suppliment Intake, Exercise Habits, Sleep Time) of 20 Elderly People

By Naomi Katayama

*Nagoya Womens University*

**Abstract-** Du to COVID-19, which started in 2020, Japanese people have continued to live in self-restraint until today. As a result, lifestyle habits have drastically changed. There are concerns about living indoors rather than going out, lack of exercise, increased alcohol consumption, irregular eating habits, and lack of sleep or excess sleep. In June 2022, we were finally able to hold a small number of health classes, and we conducted a questionnaire survey on the current living situation so that we will report the results. Participants were 20 elderly females who had been informed about the study and agreed to a consent form. Participants measured their height, weight, blood pressure, and completed a self-reported questionnaire. The content of the self-reported questionnaire survey was the smoking habit, drinking habit, exercise habit, sleeping time, and supplement intake. Participants were also asked about co-habitation.

**Keywords:** *questionnaire survey, drinking, smoking, exercise habits, sleep time.*

**GJMR-L Classification:** *DDC Code: 610.73072 LCC Code: RT81.5*



*Strictly as per the compliance and regulations of:*



© 2022. Naomi katayama. This research/review article is distributed under the terms of the Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0). You must give appropriate credit to authors and reference this article if parts of the article are reproduced in any manner. Applicable licensing terms are at <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

# Questionnaire Survey Results on Lifestyle Habits (Drinking, Smoking, Suppliant Intake, Exercise Habits, Sleep Time) of 20 Elderly People

Naomi Katayama

**Abstract-** Du to COVID-19, which started in 2020, Japanese people have continued to live in self-restraint until today. As a result, lifestyle habits have drastically changed. There are concerns about living indoors rather than going out, lack of exercise, increased alcohol consumption, irregular eating habits, and lack of sleep or excess sleep. In June 2022, we were finally able to hold a small number of health classes, and we conducted a questionnaire survey on the current living situation so that we will report the results. Participants were 20 elderly females who had been informed about the study and agreed to a consent form. Participants measured their height, weight, blood pressure, and completed a self-reported questionnaire. The content of the self-reported questionnaire survey was the smoking habit, drinking habit, exercise habit, sleeping time, and supplement intake. Participants were also asked about co-habitation. Most of the participants (60%) lived as married couples. Of the participants, 60 % were alcohol drinkers, mainly beer and wine. Sixty percent of the participants answered that they do not smoke. Among the participants, 40% used supplements, mainly used calcium (Osteoporosis prevention), Iron (anemia prevention), chlorella (nutrition supply), and green juice (vegetable shortage prevention). Most participants slept for 6 hours (30%) and 45% participants slept for 7 hours or more (45%). Of the participant, 20% had no exercise habits, and 35% took walks. As a result, many participants stayed at home for a long time due to self-restraint and did not exercise outdoors. Some participant stayed at home without even taking a walk. We would like to encourage more participants to leave their homes, walk, meet people, and help maintain a regular life.

**Keywords:** questionnaire survey, drinking, smoking, exercise habits, sleep time.

## I. INTRODUCTION

Due to the COVID-19 epidemic, which has continued for three years since 2022, Japanese people are live self-restraint at home. Due to COVID-19, by staying at home, this influences the diet, exercise habits, sleep times, etc., in Japanese life. Therefore, in this study, we conducted a questionnaire

survey on the lifestyle habits of 20 elderly females in a small-group health class for the first time in three years.

## II. MATERIAL AND METHODE

Participants signed and sealed a research consent form after receiving an explanation of the research. Participants sanitized their hands with alcohol, wore masks, and measured their body temperature in a ventilated room. In addition, the participants answered a self-reported questionnaire while sitting in seats separated by acrylic panels with a seating distance of 1 m or more. The items of the questionnaire survey for the participants were whether or not they lived together, whether or not they had support or nursing care, whether or not they quit smoking, whether they used alcohol, whether they took supplements, their exercise habits, and their sleep times. Participants also had their height, weight, and blood pressure measured.

This research has been approved after an ethical review at Nagoya Women's University (Hitowomochiitakennkyunikansuruiinkai No. 2020-9)

There are no companies, etc., with COI relationships in this research.

## III. RESULTS

Table 1 shows the ling conditions of the participants (the number of co-habitants and the presence or absence of support or nursing care), and Table 2 shows the participant's height, weight, and blood pressure. Most of the participants (60%) lived as married couples. Twenty percent of the participants lived alone. There were 20 female participants with an average age  $\pm$  Standard deviation of  $78.5 \pm 18.6$  years. Participant's mean blood pressure was within the normal range.

Table 1. Living situation survey results (20 elderly females)

Number of people living together	Alone	2 people	3 people	4 people	5 people	6 people
Status of support and nursing care	none	support 1	support 2	nursing care 1	nursing care 2	nursiing care 3 or higher
	19	1	0	0	0	0

**Author:** Nagoya Women's University, Faculty of Health Sciences, Department of Health and Nutrition. e-mail: naomik@nagoya-wu.ac.jp

Table 2. Body and blood pressure information of health class participants (20 elderly females)

	Age	Hight	Weight	Systolic blood pressure	Diastolic blood pressure
Average value	78.5	152.3	60	118.6	78.4
Standard diviation	18.6	25.6	19.4	19.7	7.3

Sixty percent of the participants answered that they do not smoke. Thirty-five percent of participants said they had quit smoking. Thirty-five percent of participants responded that they do not drink (Table 3). Of the participants, 60 % were alcohol drinkers, mainly drinking beer and wine (Table 4.)

Table 3 Smoking and drinking status of health class participants (20 dlderly females)

Smoking	No smoking	Stop to smoking	Smoking	No answer
	12	7	0	1
Drinking	No drinking	Stop to drinking	Drinking	No answer
	7	1	12	0

Table 4 Drinking frequency and content of 12 participants who drink alcohol (20 dlderly females)

Frequency	Every day	5~6/week	3~4/week	1~2/week	1~2/month
	4	3	1	2	2
Content	Japanese Sake	Cocktails	Beer	Wine	other
	1	1	7	3	0

Among the participants, 40% took supplements such as calcium, iron, vitamin B, vitamin C, cod liver oil, chlorella, and green juice (Table 5).

Table 5 Frequency and Type of Supplement of health class participants (20 dlderly females)

Frequency	Do not take	sometimes	every day	no answer
	11	2	6	1
Type	Ca	Fe	Vitamin B	Vitamin C
	1	1	1	1
Type	Chlorella	Cod liver oil	Green juice	Others
	1	1	1	

Of the participants, 20% had no exercise habits, and 35% took walks (Table 6).

Table 6 Exercise habits (20 dlderly females)

Amount of exercise per week	Almost never	1~2 hours	3~4 hours	more than 5 hours	No answer
	4	6	6	3	1
Amount of exercise per one year	Almost never	Sometimes	once a week	2 times a week	No answer
	4	2	8	5	1
Frequency of walks	Almost never	1 time per week	2~3 times per week	every day	No answer
	6	3	3	7	1

Among the participants, 30% had 6 hours of sleep, followed by 20% with 8 hours and 20% with 7 hours (Table 7).

Table 7 Sleep hours, Bedtime, and Wake up time (20 elderly females)

Sleep hours	4 hours	5 hours	6 hours	7 hours	8 hours	No answer
	2	2	6	4	5	1
Bedtime	21 o'clock	22 o'clock	23 o'clock	24 o'clock	Irregular	No answer
	2	6	8	2	2	0
Wake up time	4 o'clock	5 o'clock	6 o'clock	7 o'clock	Irregular	No answer
	1	3	11	2	3	0

#### IV. DISCUSSION

In this study, the participants lived independently (few people needed care or support), smoked less (only 20%), and drink less. Drinkers were mostly beer, around 350ml.

Among the participants, 40% used supplements, mainly used calcium (Osteoporosis prevention), Iron (anemia prevention), chlorella (nutrition supply), and green juice (vegetable shortage prevention). Most participants slept for 6 hours (30%), and 45% of participants slept for 7 hours or more (45%). Most participants went to bed at 23:00 (40%) and woke up at 6:00 (55%). Many participants had a regular life time in the city. On the other hand, since 10-15% of the participants lead an irregular lifestyle, we believe that it is necessary to encourage early to bed and early rise and light exercise (walking, etc.) that allows exposure to sunlight during the day time. It is conceivable that by confining such participants at home, their internal clocks would go out of whack without being exposed to sunlight, and their life rhythms would become irregular due to the inability to get good quality sleep. In this survey, 20% of participants did not get enough exercise, and 30% did not even walk. In the future, to increase the number of people who exercise regularly, we would like to hold more events such as health classes while checking the situation of COVID-19. Through these results, we would like to encourage more participants to go out of their homes, walk, meet people, and help maintain a regular life.

#### V. CONCLUSIONS

A self-administered questionnaire survey in a small group of 20 elderly females in a health class revealed their lack of exercise. It is possible that the COVID-19 stay-at-home policy that has continued since 2020 has had a significant impact. In the future, while assessing the situation of COVID-19, we think that it is necessary to hold more events, recruit participants, and make efforts to get people out and restore their exercise

habits. The US Physical Activity Guidelines recommend that all adults do at least 150 minutes of moderate-to-vigorous physical activity per week. It has become clear that inactive people are more likely to become severely ill than active people when contracting the coronavirus<sup>1</sup>. Under COVID-19, as a result of measuring the number of steps using a mobile phone step counting application, it is reported that it is 3000 fewer steps per day in 1-2 weeks than before COVID-19<sup>2</sup>. But on the other hand, from the UK around COVID-19, people have met 150 minutes or more of moderate-to-vigorous physical activity<sup>3</sup>. It appears that the impact of CIVD-19 on exercise habits varies considerably form country to country. This may be differences in lifestyle habits, exercise habits, and national countermeasures in each country. From now on, we think it is necessary to obtain reports from many countries and cities and compare them.

#### ACKNOWLEDGEMENTS

This study was supported by the Japanese Society of Taste Technology, 2022 and this research was partially supported by the research aid of Choju-iryō-kenkyū-kaihatsuhi, 2022 (30-14, Hirokazu Suzuki)

#### REFERENCES RÉFÉRENCES REFERENCIAS

1. Sallis R, et al. Br J Sports Med. 2021; 0: 1–8. doi: 10.1136/bjsports-2021-104080.
2. Tison GH et al. Worldwide Effect of COVID-19 on Physical Activity: A Descriptive Study. Ann Intern Med. 2020; M20-2665.
3. Smith L et al, BMJ Open Sport & Exercise Medicine, 2020; 6: e000850.